

On The Relationship Between Age and Dynamics in Elliptical Galaxies

Duncan A. Forbes, Trevor J. Ponman

School of Physics and Astronomy, University of Birmingham, Birmingham, B15 2TT

Email: forbes, tjp@star.sr.bham.ac.uk

ABSTRACT

Galaxy age estimates (mostly from spectroscopy of the central regions) are now available for many early type galaxies. In a previous paper [Forbes, Ponman & Brown 1998, ApJ, 508, L43], we showed that the offset of galaxies from the fundamental plane depends on galaxy age.

Here, using the same sample of 88 galaxies, we examine the scatter about the Faber–Jackson (F–J) relation, and find that a galaxy’s position relative to this relation depends on its age. In particular, younger ellipticals are systematically brighter in M_B and/or have a lower central velocity dispersion (σ_o). The mean relation corresponds to galaxies that are ~ 10 Gyr old.

We attempt to reproduce the observed trend of the F–J residuals with age using two simple models. The first assumes that galaxy age is tracing the last major star formation event in an elliptical galaxy. We assume that this starburst was instantaneous, centrally located and involved 10% of the galaxy by mass. The fading of this burst changes the M_B component of the F–J residuals, with time. Such a model was very successful at reproducing the B–V and Mg_2 evolution reported in our previous paper, but is unable to reproduce the strength of the F–J trend. A second model is required to describe age–correlated changes in galaxy dynamics. Following expectations from cosmological simulations, we assume that σ_o , for a galaxy of a given mass, scales with the epoch of galaxy formation, i.e. with the mean density of the Universe. Hence recently formed ellipticals have systematically lower velocity dispersions than old ellipticals. We find that a combination of these two models provides a good match to the change in F–J residuals with galaxy age. This suggests that young ellipticals will have subtly different dynamical properties to old ellipticals.

We also find that there is *not* a strong relationship between a galaxy’s age and its luminosity for our sample. This suggests that the *tilt* of the fundamental plane is not totally driven by age.

Subject headings: galaxies: elliptical, galaxies: kinematics and dynamics, galaxies: photometry

1. Introduction

Hierarchical Clustering and Merging (HCM) is the generally accepted paradigm governing the formation and evolution of elliptical galaxies. In this paradigm, elliptical galaxies are formed by the major merger of disk systems and their associated dark matter halos. This scenario for forming ellipticals was suggested in the seminal work of Toomre & Toomre (1972), and is often referred to as the ‘merger hypothesis’.

This hypothesis has gained much support over the years (see review by Barnes & Hernquist 1992). Simulations can now be used to track the stellar, gaseous and dark matter evolution of a merger from initial contact, through coalescence of the two nuclei, to the resulting remnant – an elliptical galaxy. On the observational side, this simple evolutionary sequence is more difficult to follow. The main difficulty is one of correctly identifying galaxies at different stages of the evolutionary sequence.

For the first 1–3 Gyrs after a merger the tidal tails produce a clear signature of a disk–disk merger, and acts as a ‘clock’ for age–dating the system. Examples of post–tail merger remnants are harder to identify. Indeed the lack of clear candidate protoellipticals has been called the ‘King gap’ after I. King, who pointed out this potential missing link in 1977. Balmer lines, associated with the merger–induced starburst, offer another clock. However after ~ 3 Gyr they have weakened to the point that metallicity effects are comparable to age in dictating the strength of the line. This is the so–called age–metallicity degeneracy. This degeneracy has meant that ellipticals could not be uniquely age–dated from stellar spectra. However, recently new spectroscopic observations and models have broken this degeneracy (e.g. Worthey 1994; Trager et al. 1999). The number of ellipticals with high quality nuclear spectral indices on the Lick/IDS system has grown rapidly in the last year. Combined with evolutionary model tracks, these indices can be used to estimate the age of the last major starburst in elliptical galaxies.

In an earlier paper (Forbes et al. 1998, Paper I) we used such age estimates for 88 early type galaxies to investigate the scatter about the fundamental plane. As recently as 1996, Jørgensen et al. (1996) found no dependence of the FP residuals on galaxy ellipticity, isophotal shape or relative disk contribution. They concluded that “It remains unknown what the source of [the FP] scatter is”. We showed that the

FP residuals correlate strongly with galaxy age. In addition, the deviation of a galaxy from the mean scaling relations of B–V colour and Mg_2 line index with mass, also correlates with age. The data are consistent with the steady reddening, increasing line strength and global fading of an ageing burst of star formation. This reinforces the view that galaxy age estimates are tracing the last major burst of star formation, which has been induced by a gaseous merger event.

Here we focus on dynamical effects in evolving elliptical galaxies. In particular we examine the location of ellipticals with age estimates with respect to the Faber–Jackson scaling relation.

2. A Sample of Galaxies with Age Estimates

In Table 1 we give the basic properties of the 88 early type galaxies which have age estimates available from the literature. We have included all galaxies from the study of Kuntschner & Davies (1998) in the Table, but we have excluded NGC 1399 from our analysis as it is a cD galaxy. From the large compilation of Prugniel & Simien (1996), we list the galaxy name, distance (which includes Virgocentric and Great–Attractor correction terms for $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$), total B band magnitude, central velocity dispersion and residual from the fundamental plane, i.e. $R(\sigma_o, M_B, \mu_e) = 2\log(\sigma_o) + 0.286M_B + 0.2\mu_e - 3.101$. For the two galaxies which are not part of the Prugniel & Simien list (i.e. the elliptical NGC 6127 and the S0 NGC 507), we used Faber et al. (1989) for their properties. Galaxies in our sample have a typical $M_B \sim -20.5$ and distance of $\sim 30 \text{ Mpc}$, with 3/4 in low density environments and 1/4 located in clusters.

If available, age estimates come from recent spectroscopic estimates from Trager et al. (1999), Kuntschner & Davies (1998), Tantalo et al. (1998) and Mehlert et al. (1997). Where these were not available we used the youngest ‘morphological’ age estimate from Schweizer & Seitzer (1992). The reference for each age estimate is also given in Table 1.

These age estimates require some comment. Firstly, the spectroscopic estimates come from the central regions and are luminosity weighted, so a centrally located starburst will dominate the spectral lines and hence the age estimate. Trager (1997) has shown that even a 10% by mass young starburst occurring at the galaxy centre dominates over light of the old stellar

population. Thus the age estimates, in the ideal situation of two distinct stellar populations, will largely reflect that of the young starburst. Secondly, the morphological ages have been ‘calibrated’ to a starburst model by Schweizer & Seitzer (1992). In this sense, they should also reflect the time since the last major starburst. If we separate our sample into spectroscopic and morphological ages, both subsamples reveal the trend found in paper I, i.e. FP residuals correlate with galaxy age.

In order to include some younger systems, we have included data for four good candidates for ~ 1 Gyr remnants of a major gaseous merger. In Table 2 we list their galaxy names, distances, absolute B magnitudes, velocity dispersions, FP residuals, age estimates and reference sources.

3. Galaxy Age and Luminosity

In Paper I we showed that the *scatter* about the FP was correlated with galaxy age. It is also important to determine whether the *tilt* of the FP is also strongly affected by age. If the FP tilt results in part from age effects then we might expect to see a relationship between the age of a galaxy and its luminosity.

The initial Lick sample of about 40 galaxies, suggested that low luminosity ellipticals were systematically younger than their high luminosity counterparts (Faber et al. 1995). This is opposite to the trend expected from HCM models (Kauffmann & Charlot 1998; Baugh et al. 1998). In Fig. 1 we show the absolute B band magnitude versus galaxy age for our sample. We also show the mean age and the standard deviation of the data within bins of 1 magnitude. This reveals a weak trend for increasing mean age with increasing luminosity, but given the large scatter our sample is consistent with no age trend. One interesting feature is that only giant ellipticals are very old.

Can this weak trend explain the tilt of the FP ? Prugniel & Simien (1996) found that the variation of mass-to-light ratio along the FP needed to explain the tilt required $M/L_B \propto L_B^{0.27}$. For our data set, the factor of 40 change in L_B (i.e. 4 mags) requires a change in M/L_B by a factor of 2.7 to fully account for the tilt. Using the Bruzual & Charlot (1993) models we can estimate the change in global M/L_B for a 10% by mass starburst embedded in an old population (i.e. as used in Paper I). From a mean starburst age of 5.8 Gyrs (for the low luminosity galaxies) to 7.6 Gyrs

(for the high luminosity galaxies), the M/L_B ratio changes by a factor of 1.03. Thus age variations may be responsible for a trend in M/L_B along the FP of this factor, which is much less than the factor of 2.7 change needed to fully explain the FP tilt. We conclude that the tilt of the FP, unlike the scatter about the FP, is not driven by galaxy age.

4. The Evolution of Merger-Remnants

Galaxy scaling relations such as B–V or Mg_2 vs M_B or $\log \sigma_o$ are largely sequences of metallicity vs galaxy mass (e.g. Kodama et al. 1998). In Paper I we showed that the scatter about these relations correlate with galaxy age. In particular, after a merger-induced starburst the system fades, reddens and has an increasing Mg_2 line index that is consistent with an ageing burst of star formation. This simple model provided a reasonable match to the age-related trends in these scaling relation residuals without the need for any change in the central velocity dispersion (i.e. dynamical evolution) following the merger event. Ideally we would like to predict the evolutionary changes within the full fundamental plane, however the difficulty lies in predicting the change of μ_e (or r_e) with time. This would require specific assumptions regarding the radial surface brightness profile of the old stellar population and how the central starburst size changes, and fades, with respect to it. Future simulations may indeed be able to address this issue. In this paper, we examine the overall luminosity evolution of a fading starburst combined with dynamical effects.

The Faber–Jackson relation (Faber & Jackson 1976) is another 2D scaling relation that relates absolute magnitude (or luminosity) to central velocity dispersion. It is close to an edge-on view of the FP. It was also one of the first scaling relations discovered for elliptical galaxies and was used by Lake & Dressler (1986) to study the dynamical properties of merger remnants. The best fit to 229 early type galaxies with $M_B < -18$ from the compilation of Prugniel & Simien (1996) gives

$$\log \sigma_o = 0.102(-M_B) + 0.243.$$

In Fig. 2 we show the residuals about this fit, i.e. $R(\sigma_o, M_B) = \log \sigma_o - 0.102(-M_B) - 0.243$, against galaxy age for our sample of 88 early type galaxies and the 4 candidate merger-remnants. The mean residual value for our sample is $R(\sigma_o, M_B) = -0.01$ indicating that, on average, it is well represented by the relation derived from the larger Prugniel & Simien

sample. Overlaid on the data are two model evolutionary tracks and the combination of the two. The short dashed line is the track for the model used in Paper I, i.e. a 10% (by mass) central starburst, of solar metallicity, superposed on an non-evolving old stellar population. Such a model, of a fading central starburst, provided a good representation of the residuals for the four scaling relations given in Paper I. This is not the case for the Faber–Jackson residuals, where young ellipticals generally lie below the model line and old ellipticals lie above it. A stronger burst, e.g. one involving 50% of the galaxy’s mass, would provide a steeper curve, which would give a better fit to the data. However, such a strong burst would give much stronger evolution in colour and line strength than is observed (see Paper I), as well as being improbably large when compared to observed starbursts. The burst model in Paper I assumed solar metallicity. Burst models with non-solar metallicities have only a minor impact on the Faber–Jackson track, and therefore also fail to reproduce the trend in the data.

If post-merger fading cannot explain the magnitude of the trend in the Faber–Jackson residuals with age, then this suggests that we should examine the role of dynamics. All burst models used thus far, have assumed no change in central velocity dispersion with age, for a galaxy of given mass. We now relax this assumption, and explore the consequences.

4.1. Dynamical Trends

We have seen above that fading of a central starburst, which can explain the evolution of colour and Mg_2 index rather well (see Paper I), is able to account for only about one third of the $\Delta R(\sigma_o, M_B) \approx 0.3$ dex trend in Faber–Jackson residuals for galaxies older than 1 Gyr, as seen in Fig.2. To explain the bulk of this trend, on the basis of changes in σ_o , would require a systematic change in σ_o of approximately 60% across the plot. This could be achieved in one of two ways: either (a) the central velocity dispersion increases with time after a merger in an individual elliptical galaxy, or (b) old ellipticals (i.e. those formed at high redshift) formed with systematically higher velocity dispersions (for a given mass) than their younger counterparts.

Substantial intrinsic dynamical evolution in post-merger ellipticals, over the long timescales required appears extremely unlikely. Simulations of galaxy merging (e.g. Navarro 1990) indicate that the veloc-

ity dispersion within the effective radius relaxes on a timescale < 1 Gyr after nuclear merger. Infall of tidal tail material over the next few Gyrs (Hibbard & Mihos 1995) cannot extend this timescale greatly, and in any case will have little impact on the central velocity dispersion.

Environmental effects could certainly operate over a longer timescale. Most ellipticals reside in groups and clusters, and may therefore be subject to a variety of external influences. Accretion of satellite galaxies might cause long term changes in σ_o without resetting the post-merger age, but this should act in the direction of *reducing* σ_o , relative to the mean Faber–Jackson relation (Hernquist, Spergel & Heyl 1993).

Another possibility is that σ_o does not evolve, but that ellipticals formed earlier have higher σ_o values. The HCM models predict that structures collapsing at earlier epochs will have a higher mean density (e.g. Navarro, Frenk & White 1997), and hence from the virial theorem, will have higher velocity dispersion for a given mass. A number of numerical and theoretical studies (e.g. Cole & Lacey 1996; Navarro, Frenk & White 1997; Tormen, Bouchet & White 1997; Salvador-Sole, Solanes & Manrique 1998) suggest that it is reasonable to regard the evolution of dark matter-dominated halos as proceeding via a combination of merging and accretion. The mean density of a halo, which is set at the time of a substantial merger event, is a multiple of the mean density of the Universe at that epoch. Subsequent accretion of material onto the halo does not disrupt the density distribution in its inner regions, so that the central velocity dispersion will remain essentially constant, until it is reset by a further major merger event.

Since the dark halos formed in numerical simulations have a universal form (Navarro, Frenk & White 1997), which seems to accord well with observations, the velocity dispersion will, by the virial theorem, scale in a simple way with system mass and mean density:

$$\sigma^2 \propto M/R \propto M^{2/3} \rho^{1/3} \propto M^{2/3} (1 + z_f),$$

where M and R are the mass and virial radius of the system, and ρ is its mean density, which scales (Lacey & Cole 1993) with the critical density of the Universe at the epoch (z_f) of its last major merger event. Hence a galaxy of a given luminosity (and therefore mass) will have a velocity dispersion which depends upon its ‘formation’ epoch as $\sigma \propto (1 + z_f)^{1/2}$. Assuming that the 10% starburst coincides with the fi-

nal merger which assembled the galaxy, the measured post-merger ages provide us with the value of z_f required to evaluate the strength of the velocity dispersion trend predicted by this equation. The resulting curve relating age to velocity dispersion is cosmology-dependent, via the relationship between look-back time and redshift.

As an example of the magnitude of the effects which are obtained, we have overlaid in Fig. 2 (long dashed line) the variation in Faber–Jackson residuals which result from a flat Λ CDM Universe with $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$ for $H_0 = 60 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The model has been normalised to zero residual at 10 Gyr, as Paper I showed that ellipticals lie on the fundamental plane if they have an age of ~ 10 Gyr. Changing the Hubble constant has a substantial effect on the model track, e.g. for $H_0 = 75$ the track lies about 0.1 dex lower around 1 Gyr, and is slightly steeper than the $H_0 = 60$ track for ages older than 10 Gyr. The dynamical trends match the overall magnitude of the residuals fairly well. But from paper I, we know that there is also an evolutionary effect on M_B from the fading starburst. This is shown in Fig. 2 by the short dashed line. The combination of the two effects (i.e. dynamical trends plus a burst model) are given by a solid line, which does a reasonable job of describing the trend of F–J residuals with age.

In order for this dynamical effect to be acceptable as an explanation for the bulk of the Faber–Jackson residuals, it is essential that it should not conflict with the success of the burst model in explaining the correlations involving colour and Mg_2 index, which were discussed in Paper I. In practice, the coefficients of $\log \sigma_o$ in the relations with B–V and Mg_2 explored in the Paper I are small (~ 0.15), so that 60% changes in $\log \sigma_o$ result in only minor perturbations to the successful model fits obtained with the fading burst model under the assumption of no dynamical trends.

5. Concluding Remarks

Our main result concerns the scatter about the Faber–Jackson (F–J) relation for elliptical galaxies. We found that the scatter about this scaling relation is correlated with the post-merger age of the galaxy. Young ellipticals are systematically brighter in the B band and/or have a lower central velocity dispersion than the mean relation. The mean relation corresponds to galaxies that are ~ 10 Gyr old.

We have also shown that we can reproduce the ob-

served trend of the F–J residuals with age using two simple concepts. First, we assume that the galaxy has undergone an instantaneous, centrally located starburst involving 10% (by mass) of the galaxy. Such a model was very successful at reproducing the B–V and Mg_2 evolution reported in our earlier paper, but we find that it can not adequately explain the F–J residuals with time. We propose that the bulk of the trend in F–J residuals results from a systematic variation in the internal dynamics. In particular, the central velocity dispersion of a newly formed elliptical scales with the mean density of the Universe. Such a change in mean density moves a galaxy around on the fundamental plane, but in the F–J projection of the plane it moves a galaxy away from the mean relation. The F–J offsets therefore contain information about the epoch of elliptical galaxy assembly, which may be some time after the bulk of its stars were actually formed. The strength of the dynamical trend observed favours a low value of the Hubble constant (i.e. $\leq 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$). We find that a combination of this effect and starburst fading provides a good match to the change in F–J residuals with galaxy age. The success of this model suggests that the derived starburst ages measure the time since the galaxy we see was assembled in essentially its present form. This in turn confirms the important role of recent dissipational mergers in the formation of elliptical galaxies.

Our results lend support to the view that the internal dynamical properties of an elliptical galaxy depend on when the galaxy was formed. The trend in F–J residuals we observe could have been predicted on the basis of HCM models, but as far as we are aware, they have not been explicitly mentioned in the literature. It follows that, if the merger of two spirals indeed produces an elliptical galaxy, then today’s ongoing mergers will form an elliptical galaxy subtly different from old ellipticals found in nearby clusters. We might also expect studies of the fundamental plane at redshifts $z \sim 1$ to begin to reveal dynamical differences to local ellipticals.

Acknowledgments

We thank R. Brown, P. James and A. Terlevich for help and useful discussions. We also thank S. Trager for allowing us to use his results prior to publication, and the referee (C. Lacey) for several suggestions which have improved the paper.

References

- Barnes, J. E., & Hernquist, L. 1992, *ARA&A*, 30, 705
- Baugh, C. M., Cole, S., Frenk, C. S., Lacey, C. G. 1998, *ApJ*, 498, 504
- Bica, E., & Allion, D. 1987, *A&AS*, 70, 281
- Cole, S., & Lacey, C. 1996, *MNRAS*, 281, 716
- Faber, S. M., & Jackson, R. E. 1976, *ApJ*, 204, 668
- Faber, S. M., et al. 1989, *ApJS*, 69, 763
- Faber, S. M., Trager, S., Gonzalez, J., Worthey, G. 1995, in *Stellar Populations*, eds P. C. van der Kruit and G. Gilmore (Dordrecht: Kluwer), 249
- Forbes, D. A., Ponman, T. J., & Brown, R. J. N. 1998, *ApJ*, 508, L43 (Paper I)
- Hernquist, L., Spergel, D. N., & Heyl, J. S. 1993, *ApJ*, 416, 415
- Hibbard, J. E., & Mihos, J. C. 1995, *AJ*, 110, 140
- Jørgensen, I., Franx, M., & Kjaergaard, P. 1996, *MNRAS*, 280, 167
- Kauffmann, G., & Charlot, S. 1998, *MNRAS*, 294, 705
- Kodama, T., Arimoto, N., Barger, A. J., Aragon-Salamanca, A. 1998, *A&A*, 334, 99
- Kuntschner, H., & Davies, R. L. 1998, *MNRAS*, 295, 29
- Lacey, C. & Cole, S. 1993, *MNRAS*, 262, 627
- Lake, G., & Dressler, A. 1986, *ApJ*, 310, 605
- Mehlert, D., Bender, R., Saglia, R. P., & Wegner, G. 1997, *astro-ph/9709295*
- Navarro, J. F. 1990, *MNRAS*, 242, 311
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *ApJ*, 490, 493
- Prugniel, P., & Simien, F. 1996, *A&A*, 309, 749
- Salvador-Sole, E., Solanes, J. M., & Manrique, A. 1998, *ApJ*, 499, 542
- Schweizer, F., & Seitzer, P. 1992, 104, 1039
- Tantalo, R., Chiosi, C., & Bressan, A. 1998, *A&A*, 333, 419
- Toomre, A., & Toomre, J. 1972, *ApJ*, 178, 623
- Tormen, G., Bouchet, F. R., & White, S. D. M. 1997, *MNRAS*, 286, 865
- Trager, S. C. 1997, Ph.D Thesis, University of California, Santa Cruz
- Trager, S. C., Faber, S. M., González, J. J., & Worthey, G. 1999, in preparation
- Whitmore, B. C., Miller, B. W., Schweizer, F., & Fall, S. M. 1997, *AJ*, 114, 1797
- Worthey, G. 1994, *ApJS*, 95, 107




Fig. 1.— Absolute B band magnitude versus galaxy age. Ellipticals are represented by filled symbols, S0s by open circles with a horizontal line. The mean age and standard deviation of the data within 1 mag. bins is represented by the open squares and horizontal lines. Our sample shows little, if any, trend of galaxy luminosity with age suggesting that the FP tilt is *not* strongly dependent on galaxy age.




Fig. 2.— Residual from the Faber–Jackson relation versus galaxy age. The same symbols are used as in Figure 1, with late stage mergers shown by stars. The position of NGC 3610, with $R(\sigma_o, M_B) = -0.66$, is not shown on the plot. The short dashed line shows the evolutionary track for a galaxy which has undergone an instantaneous, solar metallicity starburst involving 10% of the total galaxy mass with no change in velocity dispersion. The long dashed line shows the dynamical effects predicted for a $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ and $H_o = 60$ Λ CDM cosmology. The solid line shows the combined effects of the two dashed lines. See text for details. The evolution from late stage mergers to old ellipticals is generally consistent with expected changes in stellar population following a merger–induced starburst, including dynamical effects due to different conditions at the epoch of formation.

Galaxy (Name)	Distance (Mpc)	$-M_B$ (mag)	σ_o (km/s)	Res. (dex)	Ref.	Galaxy (Name)	Distance (Mpc)	$-M_B$ (mag)	σ_o (km/s)	Res. (dex)	Ref.
IC2006	18.11	19.23	126	-0.08	1	N4697	15.13	20.90	180	-0.28	2
IC4051	88.30	20.82	237	0.20	1	N4816	88.30	21.15	225	0.05	5
N0315	58.88	22.20	299	-0.07	2	N4915	43.85	20.22	209	-0.30	3
N0547	63.38	21.84	240	0.12	2	N5018	30.19	20.96	212	-0.40	3
N0584	63.38	20.42	223	-0.19	2	N5322	27.79	21.20	231	-0.26	3
N0596	22.28	20.05	164	-0.23	3	N5557	43.25	21.30	259	-0.15	3
N0636	22.28	19.54	166	-0.10	2	N5576	21.68	19.93	188	-0.22	3
N0720	20.79	20.47	244	0.05	2	N5638	21.68	19.76	166	0.04	2
N0821	20.98	19.92	207	0.15	2	N5812	24.54	20.01	214	-0.06	2
N1052	17.70	19.98	217	0.07	3	N5813	22.90	20.51	230	0.19	2
N1336	18.11	18.19	115	0.32	1	N5831	22.90	19.60	169	0.02	2
N1339	18.11	18.85	179	0.17	1	N5846	22.90	21.17	246	0.08	2
N1351	18.11	18.99	147	0.05	1	N5982	37.49	20.85	251	-0.10	3
N1374	18.11	19.47	213	0.24	1	N6127	64.40	21.12	226	–	2
N1379	18.11	19.76	128	-0.14	1	N6702	57.01	20.85	176	-0.20	2
N1399	18.11	20.84	338	0.17	1	N6703	29.92	20.46	173	-0.27	2
N1404	18.11	20.40	242	-0.17	1	N7454	24.32	19.40	113	-0.25	2
N1419	18.11	17.87	129	0.09	1	N7562	39.99	20.67	277	0.10	2
N1427	18.11	19.54	171	0.04	1	N7619	39.99	20.97	321	0.18	2
N1453	49.20	21.26	289	0.01	2	N7626	39.99	21.05	268	0.11	2
N1600	59.97	22.10	343	0.05	2	N7785	43.85	20.82	253	-0.02	2
N1700	50.58	21.52	243	-0.37	2						
N2300	27.66	20.56	252	0.13	2	N0474	26.79	19.64	169	0.03	3
N2778	29.24	19.03	168	0.18	2	N0507	68.20	21.98	366	–	2
N2974	28.31	20.77	211	-0.17	3	N1316	18.11	21.86	250	-0.36	1
N3193	21.57	19.90	193	-0.06	3	N1375	18.11	18.47	76	-0.23	1
N3377	10.00	18.82	146	-0.04	2	N1380	18.11	20.60	244	0.02	1
N3379	10.00	19.69	217	0.00	2	N1380A	18.11	18.65	80	0.00	1
N3605	19.76	18.20	101	-0.05	3	N1381	18.11	19.01	163	-0.03	1
N3608	19.76	19.80	201	0.12	2	N2549	17.86	19.26	159	-0.11	3
N3610	27.28	20.53	162	-0.66	3	N2685	15.84	19.02	103	-0.28	3
N3640	22.90	20.61	184	-0.29	3	N2732	27.28	19.49	170	-0.10	3
N3818	21.47	19.03	199	0.24	2	N2768	20.89	20.95	197	-0.11	3
N4125	25.94	21.53	234	-0.23	3	N2911	41.30	20.82	235	0.32	3
N4168	33.72	20.53	186	-0.13	3	N3226	21.58	19.83	191	0.31	3
N4261	31.47	21.29	316	0.08	2	N3489	10.00	18.95	138	-0.31	3
N4278	16.21	20.14	248	0.09	4	N3607	19.77	20.69	224	-0.07	3
N4283	16.21	18.19	103	-0.22	3	N3998	17.45	19.73	298	0.20	3
N4374	15.92	20.96	294	0.01	2	N4111	13.42	19.17	166	-0.30	3
N4472	15.92	21.77	300	-0.09	2	N4382	15.92	21.14	189	-0.38	3
N4478	15.92	19.01	144	-0.18	2	N5485	28.18	20.16	162	0.01	3
N4489	15.92	18.18	64	-0.28	2	N7332	15.27	19.17	138	-0.31	3
N4552	15.92	20.25	263	0.00	2	N7457	10.66	18.99	84	-0.13	3
N4649	15.92	21.35	342	0.08	2	N7585	39.45	21.02	226	-0.09	3
N4660	15.92	18.87	188	-0.03	3	N7600	40.17	20.45	223	0.07	3

Distances (assuming $H_o = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$), absolute B band magnitudes, velocity dispersions and fundamental plane residuals are from Prugniel & Simien (1996). The table lists 66 elliptical galaxies followed by 23 S0 galaxies. Note that NGC 1399 (a cD galaxy) is included in the Table for completeness but not included in the analysis. References for ages are: 1 = Kuntschner & Davies (1998); 2 = Trager *et al.* (1999); 3 = Schweizer & Seitzer (1992); 4 = Tantaló *et al.* (1998); 5 = Mehlert *et al.* (1997).

Table 2. Merger–Remnant Candidates

Galaxy (Name)	Distance (Mpc)	$-M_B$ (mag)	σ_o (km/s)	FP Residual (dex)	Age (Gyr)	Ref.
N2865	36.48	20.72	177	-0.52	1.2	1
N3156	19.95	18.81	86	-0.28	0.9	2
N3921	72.78	21.37	200	-0.26	0.8	3
N7252	52.48	21.17	177	-0.50	0.7	4

Notes to Table 2:

Distances (assuming $H_o = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$), absolute B band magnitudes, velocity dispersions, residuals from the fundamental plane are from Prugniel & Simien (1996). Galaxy ages are estimates of the time since the merger event. References for ages are: 1 = Bica & Alloin (1987); 2 = Schweizer & Seitzer (1992); 3 = Schweizer *et al.* (1996); 4 = Whitmore *et al.* (1997).